

**POISON**

Keep out of reach of children.

Read safety directions before opening or using.

**FarmSafe<sup>TM</sup>**  
**weed *TERMINATOR***

Concentrate Herbicide

For the control of weeds and grasses in non-crop areas in horticulture,  
orchards, vineyards around buildings and firebreaks.

GROUP **Z** HERBICIDE

ACTIVE CONSTITUENT: 900g/L Acetic Acid, 10g/L Hydrochloric Acid

**200 Litre**

# FarmSafe™

## weed TERMINATOR

**DIRECTIONS FOR USE** For optimum results Contact Organics FarmSafe Weed Terminator Concentrate Herbicide is intended to be used in conjunction with Contact Organics Boost. If not using Contact Organics Boost, the higher rate specified in the table below must be used for effective weed control.

**DO NOT** apply if rain is expected within 2 hours. Rainfall within 2 hours of application may wash the chemical off the foliage and repeat treatment may be required.

**DO NOT** apply if heavy rains or storms are forecast within 3 days. **DO NOT** apply by aircraft. **DO NOT** irrigate to the point of runoff for at least 3 days after application. **DO NOT** broadcast application over the entire area to be treated.

| Situation  | Weeds Controlled  | Rate  | Critical Comments  |
|--|---|---|--|
| Under trees and vines in orchards and vineyards; Non-crop areas such as verges, paths, driveways, around sheds and other buildings; firebreaks; spot spraying in lawns and turf. | Annual Weeds: Broadleaf weeds (e.g. Blackberry nightshade, Capeweed, Burr Medic), Creeping oxalis, Milk Thistle, Spear Thistle, Wireweed<br>Perennial Weeds: Broadleaf weeds (e.g. Flatweed, Hair Hawkbit, Lamb's tongue, Dandelion, Evening primrose), Couch grass, Paspalum, Perennial ryegrass, Kikuyu | 3.2L/100L of spray solution.<br>6.4L/100L of spray solution, if not using Contact Organics FarmSafe Boost | Apply to actively growing plants less than 30cm high. Spray to ensure complete coverage of unwanted plants to the point of run-off. Reapplication at around 14 days may be necessary if regrowth occurs. Perennial weeds and grasses may require repeated application to obtain long-term control. Optimal results are achieved when Contact Organics FarmSafe Weed Terminator Concentrate Herbicide is applied in full sun to small, actively growing plants. |
| Resistant weed control in the above situations.  |   | 4.8L/100L of spray solution.<br>9.6L/100L of spray solution, if not using Contact Organics FarmSafe Boost | When spraying under trees in orchards and vineyards, <b>DO NOT</b> broadcast application over the entire area to be treated. Limit treatment to spot spraying only. This product may damage lawn and turf when used as a spot spray.   |
| Paths, tiled roofs, walls, driveways, around sheds; under trees; spot spraying in lawns and turf.  | Algae, Lichen Liverwort, Moss   | 3.2L/100L of spray solution.<br>6.4L/100L of spray solution, if not using Contact Organics FarmSafe Boost | Spray to completely cover plant surface. On hard surfaces or heavy infestations, clean off by brushing after three days and reapply, if necessary. When spraying under trees in orchards and vineyards, <b>DO NOT</b> broadcast application over the entire area to be treated. Limit treatment to spot spraying only. This product may damage lawn and turf when used as a spot spray.  |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**GENERAL INSTRUCTIONS:** For optimum results Contact Organics FarmSafe Weed Terminator Herbicide should be used in conjunction with Contact Organics FarmSafe Boost. If not using Contact Organics FarmSafe Boost, use Contact Organics FarmSafe Weed Terminator Herbicide at a higher rate in accordance with the Directions for Use table. Apply to actively growing plants less than 30cm high. Spray to ensure complete coverage of unwanted plants to the point of run-off. Optimum results are obtained when Contact Organics FarmSafe Weed Terminator Concentrate Herbicide is applied in conjunction with FarmSafe Boost in full sun to small, actively growing plants. The recommended application rate for actively growing weeds at medium weed density is 500L/ha. If weed density is high, the application rate is increased. Spot spraying is typically conducted at an application rate up to 1000L/ha. For woody weeds and resistant weeds, the concentration of the spray mixture should be increased by 50% as per the Directions for Use table. Perennial weeds and grasses may require repeated application to obtain long-term control. Contact Organics FarmSafe Weed Terminator Concentrate Herbicide works by dehydrating the plant. Results are seen as browning and wilting, usually within hours of application. To be effective, Contact Organics FarmSafe Weed Terminator Concentrate Herbicide must come into contact with the green parts of the plant. Vinegar odour dissipates within 30 minutes to 1 hour.

**MIXING INSTRUCTIONS:** Do not mix or store in steel or galvanized tanks. Mix with clean water only. **DO NOT** mix more than is needed at any one time. Use only plastic lined, fibreglass, aluminium or stainless steel containers for mixing and application. Wash out spray equipment thoroughly after use. The use of a nonionic wetting agent is recommended.

1. Half fill the spray tank with clean water and start agitation.
2. Add the required amount of Contact Organics FarmSafe Boost to the spray tank.
3. With agitator still running, add the required amount of Contact Organics

FarmSafe Weed Terminator Concentrate Herbicide and the remaining water to the tank and mix thoroughly.

4. Add wetting agent, if required, near the end of the filling process to minimise foaming.

Always maintain adequate agitation during application and use tank mix promptly. This product may be corrosive to metal parts of the sprayer. Thoroughly flush tanks, pumps, and nozzles with water after use. The use of a coarse nozzle to achieve even distribution is recommended.

**RESISTANCE WARNING:** FarmSafe Weed Terminator Concentrate Herbicide is a member of Group Z a group of herbicides with unknown and probably diverse sites of action. The product has a desiccating action on weeds. For weed resistance management, the product is a Group Z herbicide. Some naturally occurring weed biotypes resistant to the product and other Group Z herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group Z herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Contact Organics accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

**PRECAUTIONS:** Do not spray directly on humans, pets, exposed food, food utensils, or food preparation areas.

**PROTECTIONS:** PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS: **DO NOT** apply under weather conditions, or from spray equipment, that may cause spray to drift into nearby susceptible plants, crops, cropping lands or pastures. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT **DO NOT** contaminate wetlands or watercourses with this product or used containers.

**STORAGE AND DISPOSAL:** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple-rinse containers before disposal. Dispose of rinsate or any undiluted chemical according to state/territory legislative requirements. If recycling, replace cap and return

clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**SAFETY DIRECTIONS:** Corrosive. Attacks eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID INSTRUCTIONS:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766. If swallowed, do **NOT** induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. First Aid Warnings - **DO NOT** inhale vapour when handling the concentrated product



**Contact Organics FarmWeed Terminator Concentrate Herbicide**  
 APVMA Approval 86050/120802  
 TM Trademark of Contact Organics Pty Ltd  
**Contact Organics Pty Ltd**  
 PO Box 5087 Glenferrie South Victoria 3122